



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NUMBER THEORY.

218. Proposed by ELIJAH SWIFT, University of Vermont.

If p is a prime > 3 , show that $\sum_{a=1}^{p-1} \frac{1}{a^2} \equiv 0 \pmod{p}$.

II. SOLUTION BY THE PROPOSER.

It may be proved (Bachmann, *Niedere Zahlentheorie*, page 155) that $A_{p-3} = 1 \cdot 2 \cdot 3 \cdots p - 3 + 1 \cdot 3 \cdot 4 \cdots p - 2 + \cdots$, taking all possible combinations of the numbers $1, 2, \cdots p - 1, p - 3$ at a time, is divisible by p .

Then

$$A_{p-3} = (p-1)! \sum_{\substack{a=1 \\ b=1}}^{a=p-1 \\ b=p-1} \frac{1}{a \cdot b} \quad a \neq b.$$

Then

$$\sum \frac{1}{a \cdot b} \equiv 0 \pmod{p}.$$

This sum may be written

$$\frac{1}{2} \left\{ \sum_{b=1}^{p-1} \frac{1}{1 \cdot b} - \frac{1}{1^2} + \sum_{b=1}^{p-1} \frac{1}{2 \cdot b} - \frac{1}{2^2} + \cdots + \sum_{b=1}^{p-1} \frac{1}{(p-1)b} - \frac{1}{(p-1)^2} \right\}.$$

But

$$\sum_{b=1}^{p-1} \frac{1}{a \cdot b} = \sum_{b=1}^{p-1} \frac{b}{b \cdot a} = \sum_{b=1}^{p-1} b \equiv 0 \pmod{p}.$$

Hence,

$$\sum \frac{1}{ab} \equiv - \sum_{a=1}^{p-1} \frac{1}{a^2} \equiv 0 \pmod{p}.$$

QUESTIONS AND DISCUSSIONS.

EDITED BY U. G. MITCHELL, University of Kansas.

NEW QUESTIONS.

26. Why should not the nomenclature of mathematics be made uniform? For example, why call a circle a *portion of a plane* in elementary geometry and a *curve* in analytic geometry? Why call a sphere a *ball* at one time and a *surface* at another time? And so on through all the configurations of two- and three-dimensional geometry.

REPLIES.

18. In view of the present pressure for saving time and gaining efficiency, what are the most important sources of economy in the mathematical courses of the high school and the first two years in college?

REPLY BY ERNEST W. PONZER, Leland Stanford Junior University.

This question assumes the existence and application of the fundamental principles of efficiency¹ in connection with our college instruction in mathematics. No doubt the editors are aware of the fact that many college professors would hardly make this assumption and that many others would regard the problem involved as quite trivial. If such is not the case, why, for instance, in many large universities is instruction in the mathematics of the freshman year given to a majority of the students enrolled in these courses by assistants in the depart-

¹ Cf. article "A Study in Efficiency," *School Science and Mathematics*, October, 1910, pp. 579-81.